

The Effects of Auditory Integration Training (AIT) on Mismatch Negativity in Children with Autism



Tato Sokhadze¹ Kirk Akaydin¹, Stephen Edelson, ² Sally Brockett² Marie Hensley, ¹ Lynzee Alworth¹, Jeff Weihing¹, Yao Wang¹, Manuel F Casanova ¹ ¹Department of Psychiatry & Behavioral Sciences, Univerity of Louisville School of Medicine, ²Autism Research Institute, San Diego, CA

Introduction & Background

> Autism is a neurological developmental disorder present in 1 of 88 children

➤ Visible symptoms of autism include lack of social interaction, communication, and restricted range of interests.

Children with autism have atypical psychophysiological reasponses both of central and autonomic nervous system


Auditory Evoked Potentials (Mismatch Negativity)

- > Children with autism are featured by sensory over sensitivity including excessive reactivity to sounds
- ➤ Reactivity to sound can be measured using auditory evoked potentials recorded during presentation of the sounds of different frequency and recording EEG responses
- Mismatch Negativity (MMN) is one of the early auditory potentials reflecting differences of evoked potentials between rare (20%) and frequent (80%) sounds.

 MMN is typically recorded at the fronto-central EEG
- recording sites and represents a difference wave (rare-minus-frequent) occurring within 130 190 ms post-stimulus ➤ Along MMN other fronto-central evoked potentials (P2a and P3a) are related to processing auditory stimuli and
- The aims of the AIT part of the study was application of Berard's AIT techniques in children in autism and monitoring MMN, P2a, and P3a changes following AIT course.

Rationale behind Auditory Integration Training







sounds.

- Auditory integration training, as developed by French totalaryngologist Guy Berard and based on the work of his predecessor, Alfred Tomatis, typically consists of 20 half-hour sessions of listening to specially modulated music over a 10- to 20-day period. AIT has been reported to be beneficial in several conditions, including be beneficial in several conditions, including autism, ADHD, dyslexia, and hypersensitive hearing at certain frequencies.

Methods of EEG recording and AIT procedure



➤ EEG data were acquired with a 128 channel Electrical Geodesics system (v. 200) consisting of Geodesic Sensor Net consisting of Geodesic Sension Feet electrodes, Net Amps, and Net Station software (v. 4.0) [Electrical Geodesics Inc., OR]. EEG data are sampled at 500 Hz, 0.1-100 Hz analog filtered.

100 Hz analog filtered.

> All stimulus presentation and behavioral response collection were controlled by a PC running E-prime software [Psychology Software Tools, PA].

Mismatch Negativity (MMN) is one of the early auditory potentials reflecting differences of evoked notentials between rare (20%) and frequent (80%) sounds. Test take 10 min to complete

Administration of Bernard's Auditory Integration Technique (AIT)

➤_Patients were given MMN test pre and post Auditory Therapy Patients were given Audiograms are and post

➤ AIT was administered through the Earducator.

Patients listened to modulated music in association with audiogram results for 30 minute sessions, twice a day for 10 days.



Iethods: ANS Activity Measurements during AIT Audiometry and Behavioral Evaluations

Measurement of physiologic data from the therapeutic techniques

The auditory evoked potentials were recorded pre and post-AIT course using EGI EEG system.





>Comprehensive Performance Index (CPI) and Aberrant Behavior Checklist (ABC) behavior survey were administered to patients pre and post therapeutic

> In this study, we investigated evoked potentials before and after AIT

- In this study, we investigated evoked potentials before and after ATI in dildten between the ages of and 2 (1 Ne-12 an ATI near ange 13.7 ys, SD4-2). There were 2 girls in in ATI group, ys, SD4-2). There were 2 girls in in ATI group, measurements in Minimath Negativity test (menta age 149, SD+3.3). > Participants with ASD were recruited from across the U.S. through the Weiskoff Child Poslutation Center and FEAT and were evaluated by Dt. Lounie Sears to confirm autism diagnosis (DSM-IV, ADI-R)). > Participants were both high and too functioning autistic children.
- Participating subjects and their parents (or legal guardians) were provided with all information regarding the study, and the consent and assent forms approved by the IRB were reviewed and signed.

Results: 2D topo plots and ERP changes post-AIT





Aberrant Behavioral Changes after AIT



Baseline and Post AIT MMN in autism, and controls



- » Berard vAIT resulted in significant decrease of Irribbility, Byperactivy, Lehapsescore, and improved and Legority Campan, and Receptive Language Sectors on CPJ. > AIT improved and Legority Campan, and Receptive Language Sectors on CPJ. > AIT improved and Legority Campan, and L